

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Orthopaedic Rheumatology

Journal homepage: www.ijor.org

Letter to Editor

Effect of combination of bone marrow aspiration concentrate (BMAC) and platelet-rich plasma (PRP) in anterior cruciate ligament reconstruction

Ashim Gupta^{1,2,3,4*}, Surya Prakash Sharma⁵

¹Regenerative Orthopaedics, Noida, Uttar Pradesh, India

²Future Biologics, Lawrenceville, Georgia 30043, USA

³South Texas Orthopaedic Research Institute (STORI Inc.), Laredo, Texas 78045, USA

⁴Indian Stem Cell Study Group (ISCSG) Association, Lucknow, Uttar Pradesh, India

⁵Northern Railway Central Hospital, New Delhi, Delhi, India



ARTICLE INFO

Article history:

Received 11-12-2023

Accepted 18-01-2024

Available online 13-02-2024

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

To the Editor,

We read a recently published article by Lin et al.¹ titled, “Effect of bone marrow aspiration concentrate and platelet-rich plasma combination in anterior cruciate ligament reconstruction: a randomized, prospective, double-blinded study”. In this study, the authors investigated the effect of combination of bone marrow aspiration concentrate (BMAC) and platelet-rich plasma (PRP) compared to the PRP alone and the control group (no orthobiologic augmentation) in enhancing graft maturation and tendon-bone tunnel interfacial healing post anterior cruciate ligament (ACL) reconstruction (ACLR). The results demonstrated inadequate improvements in clinical function, graft maturation and tendon-bone tunnel interfacial healing in both BMAC+PRP and PRP alone groups compared to the control group. Although it was an interesting study, we have two concerns that we would like to communicate.

First, the authors did not include BMAC alone group as an active comparator. There are insufficient studies assessing the role of BMAC in graft maturation and/or enhancing ligamentization during ACL reconstruction. In a recent study, Forsythe et al.² demonstrated that

BMAC augmentation has the potential to accelerate the ligamentization of allografts utilized during ACLR, however no significant differences were observed between the BMAC and the control groups in terms of attaining minimally clinically important difference. We believe inclusion of BMAC alone group would have added valuable insights into the limited currently available literature on this topic.

Second, it is imperative that orthobiologics used, i.e., BMAC and PRP, must be characterized. For PRP, it is important to report the platelet count, leukocyte concentration, content of red blood cells (RBCs) and use of exogenous activator(s).³ Several studies have reported that a platelet concentration of 5-7 times compared to the baseline whole blood is essential to promote cell proliferation, mesenchymal stem/stromal cells recruitment and tissue healing.^{3,4} A recent editorial summarized the harmful effects of RBCs and recommended it to be avoided in PRP formulations.⁵ Interestingly, a recent pre-clinical study also demonstrated superiority of RBCs depleted BMAC over regular BMAC.⁶ In addition, for BMAC, it is important to report the content of total nucleated cells, expression of hematopoietic and mesenchymal stem cells markers, colony forming units, and platelet counts.⁷ The lack of standardized formulation protocols and patient variables, including age,

* Corresponding author.

E-mail address: ashim6786@gmail.com (A. Gupta).

comorbidities, concomitant medications taken, etc., have been attributed to subpar outcomes associated with studies utilizing orthobiologics, and all of this information is critical to ensure repeatability and reproducibility of studies outcomes.^{8–10}

List of Abbreviations

BMAC-Bone Marrow Aspiration Concentrate, PRP-Platelet-rich Plasma, ACL-Anterior Cruciate Ligament, ACLR-Anterior Cruciate Ligament Reconstruction, RBCs-Red Blood Cells.

Conflict of Interest

The authors declare that they have no competing interests.

Source of Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Authors' Contributions

AG conceptualized the study and wrote the initial manuscript draft. AG and SPS reviewed and edited the manuscript draft and have read and agreed to the published version of the manuscript.

Acknowledgements

None.


References

1. Lin YC, Chen YJ, Fan TS, Chou PH, Lu CC. Effect of bone marrow aspiration concentrate and platelet-rich plasma combination in anterior cruciate ligament reconstruction: a randomized, prospective, double-blinded study. *J Orthop Surg Res.* 2024;19(1):4. doi:10.1186/s13018-023-04512-y.
2. Forsythe B, Chahla J, Korrapati A, Lavoie-Gagne O, Forlenza E, Diaz CC, et al. Bone Marrow Aspirate Concentrate Augmentation May Accelerate Allograft Ligamentization in Anterior Cruciate Ligament

Reconstruction: A Double-Blinded Randomized Controlled Trial. *Arthroscopy.* 2022;38(7):2255–64.

3. Mautner K, Malanga GA, Smith J, Shiple B, Ibrahim V, Sampson S, et al. A call for a standard classification system for future biologic research: the rationale for new PRP nomenclature. *PM R.* 2015;7(4 Suppl):53–9.
4. Giusti I, Ruggetti A, Ascenzo D, Millimaggi S, Pavan D, Dell A, et al. Identification of an optimal concentration of platelet gel for promoting angiogenesis in human endothelial cells. *Transfusion.* 2009;49(4):771–8.
5. Gupta A, Maffulli N, Jain VK. Red Blood Cells in Platelet-Rich Plasma: Avoid If at All Possible. *Biomedicines.* 2023;11(9):2425. doi:10.3390/biomedicines11092425.
6. Pezzanite LM, Timkovich AE, Sikes KJ, Chow L, Hendrickson DA, Becker JR, et al. Erythrocyte removal from bone marrow aspirate concentrate improves efficacy as intra-articular cellular therapy in a rodent osteoarthritis model. *Ann Transl Med.* 2023;11(9):311. doi:10.21037/atm-22-4256.
7. Atluri S, Murphy MB, Dragella R, Herrera J, Boachie-Adjei K, Bhati S, et al. Evaluation of the Effectiveness of Autologous Bone Marrow Mesenchymal Stem Cells in the Treatment of Chronic Low Back Pain Due to Severe Lumbar Spinal Degeneration: A 12-Month, Open-Label, Prospective Controlled Trial. *Pain Physician.* 2022;25(2):193–207.
8. Gupta A, Potty AG, Maffulli N. Allogenic platelet-rich plasma for treatment of knee and hip osteoarthritis. *Front Pain Res (Lausanne).* 2023;p. 1216190. doi:10.3389/fpain.2023.1216190.
9. Gupta A, Jeyaraman M, Maffulli N. Common Medications Which Should Be Stopped Prior to Platelet-Rich Plasma Injection. *Biomedicines.* 2022;10(9):2134. doi:10.3390/biomedicines10092134.
10. Muthu S, Jeyaraman M, Narula A, Ravi VR, Gandhi A, Khanna M, et al. Factors Influencing the Yield of Progenitor Cells in Bone Marrow Aspiration Concentrate-A Retrospective Analysis of 58 Patients. *Biomedicines.* 2023;11(3):738. doi:10.3390/biomedicines11030738.

Author biography

Ashim Gupta, Founder and President, CEO, Chief Science Officer, and Professor  <https://orcid.org/0000-0003-1224-2755>

Surya Prakash Sharma, Senior DMO

Cite this article: Gupta A, Sharma SP. Effect of combination of bone marrow aspiration concentrate (BMAC) and platelet-rich plasma (PRP) in anterior cruciate ligament reconstruction. *IP Int J Orthop Rheumatol* 2023;9(2):115-116.